



PubMed N	ucleotide Protein Genome Structure PopSet Taxonomy OMIM Books					
Search PubMed	for Go Clear					
	∠Limits Preview/Index History Clipboard Details					
<u></u>	Display Abstract ▼ Sort ▼ Save Text Clip Add Order					
Entrez	☐1: Glycoconj J 1996 Apr;13(2):315-9 Related Articles, NEW Books, LinkOut					
PubMed	Use of diethyl squarate for the coupling of oligosaccharide amines					
	to carrier proteins and characterization of the resulting neoglycoproteins by MALDI-TOF mass spectrometry.					
PubMed Services	Kamath VP, Diedrich P, Hindsgaul O.					
	Department of Chemistry, University of Alberta, Edmonton, Canada.					
Related Resources	The 8-methoxycarbonyloctyl glycosides of GlcNAc beta, Gal beta 1-4Glc beta, Fuc alpha 1-2Fuc alpha 1-3GalNAc beta and Fuc alpha 1-2Gal beta 1-3[Fuc alpha 1-4]GlcNac beta were converted to primary amines by reaction with neat ethylenediamine and then coupled to bovine serum albumin (BSA) using diethyl squarate as the connector. The average degree of incorporation of the sugar onto the protein, as well as the molecular weight distribution, could be conveniently determined using matrix assisted laser desorption ionization/time of flight (MALDI-TOF) mass spectrometry thus avoiding cumbersome structure-dependent colour-tests or analysis of cleaved ligand. The present coupling method has the advantages of proceeding under very mild conditions, yielding controlled incorporation values and can reliably be used for the coupling of very small amounts (mg) of oligosaccharide.					
	PMID: 8737256 [PubMed - indexed for MEDLINE]					
	Display Abstract Sort Save Text Clip Add Order					

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

i6So-pc-linux-gnu Mar 14 2002 16:00:17







J NCDI	I do m	400	of Medicine	NLM		
PubMed N	ucleotide Protein Genome	Structure	PopSet	Taxonomy	OMIM Books	
Search PubMed	for				o Clear	
	∠Limits Preview/Index	History	Clipboard	Details	<u> </u>	
	Display Abstract v	Sort 🔻	Save Text	Clip Add	Order	
Entrez PubMed	□1: J Immunol Methods 197	79;25(4):323-	.35 Relat	ed Articles,	NEW Books , LinkOut	
	Coupling of acid labile Salmonella specific oligosaccharides to macromolecular carriers.					
PubMed	Svenson SB, Lindberg	AA.				
Services	A coupling method for consistency isolated from S. typhimus described. Arylamine group oligosaccharides by reach subsequent conversion to derivatives saccharides word different carrier protein and elicited in rabbits bo	trium O-polysoups were into ting them with the correspondence covalent ins. The result anti-harpton	saccharide to no roduced into the continuous acceptance of the continuous continuous continuous conjugates and anti-care continuous	nacromolecume terminal responsible the terminal responsible the terminal responsible to the terminal responsibility to the terminal respons	alar carriers is reducing end of amine. After sothiocyanato sylamine groups y immunogenic in specific	
Related Resources	antibodies. Some of the advantages of this coupling procedure are: (i) it can be used with oligosaccharides containing highly acid or alkali labile structures and/or glycosidic linkages, (ii) it produces conjugates with high degrees of substitution at low saccharide/protein molar input ratios, (iii) it does not grossly affect the immunogenic specificities of the carrier protein, and (iv) it is suitable for preparation of highly substituted affinity columns, e.g., coupling to a polyacrylamide matrix. PMID: 85676 [PubMed - indexed for MEDLINE]					
	PIVIID: 830/0 [Publised - Indexed for MEDERINE]					
	Display Abstract ▼	Sort ▼	Save Text	Clip Add	Order	

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

sparc-sun-solaris2.8 Mar 21 2002 14:48:55